arXiv:1202.1619v1 [math.GN] 8 Feb 2012

Some Open Problems in Topological Algebra

Taras Banakh, Mitrofan Choban, Igor Guran, Igor Protasov

This is the list of open problems in topological algebra posed on the conference dedicated to the 20th anniversary of the Chair of Algebra and Topology of Lviv National University, that was held on 28 September 2001.

Problem 1 (Choban). *Is every topological group a quotient group of a zero-dimensional topological group of the same weight? When an almost metrizable topological group is a quotient of a zero-dimensional group of the same weight?*

A space X is zero-dimensional if $\dim X = 0$. A topological group is *almost metrizable* if contains a compact subset of countable character. Let us make some comments to this problem. If a topological group G is a quotient of an almost metrizable group H with $\mathrm{ind} H = 0$, then G contains a zero-dimensional compact subgroup of countable character in G.

If H is a zero-dimensional subgroup of countable character in a group G and $xh = hx$ for every $x \in G$ and $h \in H$, then G is a quotient group of some almost metrizable zero-dimensional group of the same weight. In particular the answer to the first part of Problem 1 is positive for metrizable groups (see [8], [6]) and for almost metrizable abelian groups. A.V.Arhangel'skii proved that every topological group is a quotient of a zero-dimensional σ-discrete group. For universal algebras this fact was proved in [7].

Problem 2 (Choban). *Under which conditions is the free universal algebra of an uncountable signature over a metrizable space X paracompact? In particular, is the free topological linear space $L(X)$ of X over the discrete field of real numbers paracompact?*

A semigroup S with the identity e endowed with a topology is called a *left* (resp. *right*) *bounded* if for every neighborhood U of e there is a finite subset F of S such that $S = FU$ (resp. $S = UF$); S is called *bounded* if S is both left and right bounded.

For a topological space X let $S(X) \subset X^X$ be the semigroup of all continuous self-mappings of X endowed with the topology of pointwise convergence (i.e., the topology inherited from the Tychonov product X^X). A topological space X is called *homogeneous* if for any points $x_1, x_2 \in X$ there is a homemorphism h of X with $h(x_1) = x_2$.

Problem 3 (Protasov). *Is the semigroup $S(X)$ left bounded for every zero-dimensional compact homogeneous space?*

The answer is positive provided X has a base of the topology, consisting of pairwise homeomorphic clopen subsets, see [11].

Problem 4 (Protasov)**.** *Is the semigroup $S(X)$ right bounded for any zero-dimensional homogeneous space X?*

The answer is positive provided X has a base of the topology, consisting of clopen subsets homeomorphic to X, see [11].

Let X be a topological space. A subgroup H of $S(X)$ is called *distal* if for any distinct points $x_1, x_2 \in X$ and any point $x \in X$ there is a neighborhood U of x such that $\{h(x_1), h(x_2)\} \not\subset U$ for all $h \in H$. It is proven in [11, Theorem 4] that a left bounded subgroup $H \subset S(X)$ is distal provided H acts transitively on X.

Problem 5 (Protasov)**.** *Let X be a compact space and H be a distal subgroup of $S(X)$, acting transitively on X. Is H left bounded?*

Under a *left-topological group* we understand a pair (G, τ) consisting of a group G and a topology τ invariant with respect to the left shifts $l_g : x \mapsto gx$. If, in addition τ is invariant with respect to the right shifts, then (G, τ) is called a *semitopological group.* A semitopological group G with continuous inverse mapping $x \mapsto x^{-1}$ is called a *quasitopological group.* If the group operation of G is continuous with respect to the topology τ, then (G, τ) is a *paratopological group.* If, additionally, the operation of taking the inverse is continuous, then (G, τ) is a topological group.

It is well known that σ-compact topological groups have countable cellularity [14] while compact topological groups support a strictly positive probability measure (i.e., a Borel probability measure μ such that $\mu(U) > 0$ for any nonempty open subset U of the group). Recently T. Banakh and O. Ravsky [3] (see also [4]) proved that any bounded paratopological group G has countable cellularity (moreover, each cardinal of uncountable cofinality is a precaliber of G). On the other hand, according to [12] for every infinite cardinal τ there is a left bounded left topological group of cellularity τ.

Problem 6 (Protasov)**.** *Let G be a bounded semitopological (quasitopological) group. Has G countable cellularity?*

To answer Problem 6 in negative it suffices to give a positive answer to

Problem 7 (Protasov)**.** *Does every zero-dimensional compact homogeneous space admit the structure of a left topological group?*[1] *In particular, does the countable power K^ω of a first-countable zero-dimensional compact space K admit the structure of a left topological group?*

Let us remark that according to Motorov Theorem the countable power K^ω of each first-countable zero-dimensional compact space K is homogeneous, see [2].

On the other hand, a positive answer to Problem 6 would follow from the positive answer of

---

[1] This problem was answered in negative by Banakh in [T.Banakh, *A homogeneous first-countable zero-dimensional compactum failing to be a left-topological group*, Mat. Stud.29 (2008) 215–217].

Problem 8 (Protasov)**.** *Does every compact left topological group G support a strictly positive probability Borel measure?*

There exists a compact left-topological group admitting no invariant probability Borel measure, see [10].

Problem 9 (Protasov)**.** *Can every non-discrete topological group G be algebraically generated by a nowhere dense subset?*[2]

Let us mention that each countable topological group is algebraically generated by some closed discrete subset (see reference in [13]) while every left topological group is algebraically generated by some subset with empty interior [13].

A paratopological group G is *left ω-bounded* (resp. *right ω-bounded*) if for any neighborhood $U \subset G$ of the origin there is a countable subset $F \subset G$ with $G = FU$ (resp. $G = UF$); G is *ω-bounded* if it is both left and right ω-bounded.

Problem 10 (Guran)**.** *Is a paratopological group G left ω-bounded if it is right ω-bounded?*

For topological groups the answers to the last two questions are in positive, see [9].

A subset A of a topological group G is called *o-bounded* if for any sequence $(U_n)_{n\in\omega}$ of neighborhoods of the origin of G there is a sequence $(F_n)_{n\in\omega}$ of finite subsets of G such that $A \subset \bigcup_{n\in\omega} F_n U_n$. It is clear that each σ-compact topological group is o-bounded while each o-bounded group is ω-bounded. Next, given a subset A of a topological group G, consider the following game OF(A) (abbreviated from Open-Finite). Two players, I and II, choose at every step $k \in \omega$ a neighborhood $U_n \subset G$ of the origin, and a finite subset F_n of G, respectively. At the end of the game, the player II is declared the winner if $A \subset \bigcup_{n\in\omega} F_n U_n$. It is easy to see that for a σ-compact group G the player II has a winning strategy in the game OF(G). On the other hand, if a topological group G is not o-bounded, then the player I has a winning strategy in OF(G).

Problem 11 (Banakh)**.** *Is there a (metrizable) o-bounded topological group G such that the player I has a winning strategy in the game OF(G)?*

Such a group, if exists, cannot be analytic and abelian (more generally, a SIN-group). We remind that a topological space X is analytic if it is a metrizable continuous image of a separable complete metric space. On the other hand, the group $\mathbb{Z}^\omega$ contains a dense o-bounded G_δ-subset A such that the first player has a winning strategy in the game OF(A), see [5].

Problem 12 (Banakh)**.** *Let A be a compact subset of the real line $\mathbb{R}$ such that $A - A = \{a - a' : a, a' \in A\}$ is a neighborhood of zero in $\mathbb{R}$. Has the sum $A + A + A + A$ non-empty interior in $\mathbb{R}$?*

---

[2] The answer to Problem 6 is “No”: under CH the topological group $\mathbb{R}^{\omega_1}$ contains a dense non-separable subgroup G such that each meager subset of G generates a metrizable separable subgroup of G; see Proposition 13.8 in [I.Protasov, T.Banakh. Ball stuctures and colorings of graphs and groups // (Mat. Stud. Monograph Series. 11), VNTL Publ. 2003, 148p.]

References

[1] *A.V.Arhangel'skii.* Any topological group is a factor group of a zero-dimensional topological group // Dokl. Akad. Nauk SSSR, 258 (1981), 1037-1040,(English transl. Soviet Math. Dokl. 23(1981)).

[2] *A.V. Arkhangelskii.* Topological homogeneity. Topological groups and their continuous images // Uspekhi Mat. Nauk. 42 (1987) 69–105, (English transl. Russian math. Surveys. 42 (1987) 83–131).

[3] *T. Banakh, O. Ravsky.* Oscillator topologies on paratopological groups and related number invariants // Algebraical Structures and their Applications, Kyiv: Inst. Mat. NANU, (2002) 140–153.

[4] *T. Banakh, O. Ravsky.* Chain conditions and invariant measures on paratopological groups (in preparation).

[5] *T. Banakh, L. Zdomsky.* Games on multicovered spaces and their application in topological algebra, in preparation.

[6] *V.K. Bel'nov.* On zero-dimensional groups // Dokl. Akad. Nauk SSSR, 226(1976), 249-252,(English transl. Soviet Math. Dokl. 17(1976)).

[7] *M.M. Choban.* On the theory of topological algebraic systems // Trans. Amer. Math. Soc. 48 (1986),115-159.

[8] *M.M. Choban.* Some questions in the theory of topological groups // General Algebra and Discrete Geometry (B.A.Shcerbakov, editor). – Shtiintsa, Kishinev, 1980, p.120-135.

[9] *I. Guran.* On topological groups close to being Lindelöf // Soviet Math. Dokl. 23 (1981), 173–175.

[10] *P. Milnor, J. Pym.* Haar measure for compact right topological groups // Proc. Amer. Math. Soc. 114 (1992) 387–393.

[11] *I. Protasov.* On totally bounded semigroups of continuous mappings (preprint).

[12] *I. Protasov.* On Souslin number of totally bounded left topological groups // Ukr. Mat. Zhurn. 54:11 (2002), 1581–1584.

[13] *I. Protasov.* Small systems of generators of groups // Mat. Zametki, 76:3 (2004), 420–426.

[14] *M. Tkachenko.* Introduction to topological groups // Topology Appl. **86** (1998), 179–231.